

Sub B1
A2

1. (Amended) A method for determining the efficiency of an amplification of a target nucleic acid comprising the steps of:
 - a) a dilution series of the target nucleic acid is prepared;
 - b) the target nucleic acid is amplified under defined reaction conditions and the amplification is measured in real-time;
 - c) a defined signal threshold value is set;
 - d) for each dilution the cycle number is determined at which the signal threshold value is exceeded; and
 - e) the amplification efficiency is determined as a function of the original amount of target nucleic acid.
2. (Amended) The method of claim 1, wherein the efficiency of an amplification is determined by:
 - a) preparing a dilution series of the target nucleic acid;
 - b) amplifying the target nucleic acid under defined reaction and the amplification of the nucleic acid being measured in real-time;
 - c) setting a defined threshold value;
 - d) determining the cycle number for each dilution at which the signal threshold value is exceeded;
 - e) determining a non-linear continuously differentiable function of a logarithm of the copy number of target nucleic acid used for the amplification as a function of the cycle number at which the signal threshold value is exceeded; and
 - f) calculating the amplification efficiency E from the function determined in step e).
3. (Amended) The method of claim 1, wherein the efficiency of an amplification is determined by:
 - a) preparing a dilution series of the target nucleic acid;
 - b) amplifying the target nucleic acid under defined reaction, the amplification of the nucleic acid being measured in real-time;
 - c) setting a defined threshold value;
 - d) determining the cycle number for each dilution at which the signal threshold value is exceeded;

- A2
Sub B1
- e) determining a non-linear continuously differentiable function of the cycle number determined in step d) as a function of the logarithm of the copy number of target nucleic acid used in each case; and
- f) calculating the amplification efficiency E from the function determined in step e).

- A3
5. (Amended) The method of claim 3, wherein the amplification efficiency E of a certain original amount of target nucleic acid is determined as the reciprocal negative local 1st derivative of the continuously differentiable function from step e).
6. (Amended) The method of claim 2, wherein the non-linear continuously differentiable function from step e) is determined with a polynomial fit.

- Sub B2
- A4
8. (Amended) A method for quantification of a target nucleic acid in a sample relative to a reference nucleic acid comprising the steps of:
- a) determination of the amplification efficiencies of the target nucleic acid and of the reference nucleic acid under defined amplification conditions as claimed in claim 1;
 - b) amplification of the target nucleic acid contained in the sample as well as of the reference nucleic acid contained in the sample under the same defined amplification conditions;
 - c) measurement of the amplification of the target nucleic acid and of the reference nucleic acid in real time; and
 - d) calculation of the original ratio of target nucleic acid and reference nucleic acid in the sample by correction of the ratio derived from step c) with the aid of the amplification efficiencies determined in step a).
9. (Amended) A method for relative quantification of a target nucleic acid relative to a reference nucleic acid and standardized with a calibrator sample comprising the steps of:
- a) preparation of a common or two separate dilution series of target nucleic acid and reference nucleic acid;

- Sub B2
- 11
- b) amplification of the various dilutions of target nucleic acid and reference nucleic acid under defined reaction conditions, the amplification of the nucleic acid being measured in real-time;
 - c) setting defined signal threshold values for the target nucleic acid and reference nucleic acid;
 - d) determining the cycle number C_p to which the signal threshold values defined for the target nucleic acid and reference nucleic acid are exceeded in each dilution;
 - e) determining a continuously differentiable function of the C_p values determined in d) as a function of the logarithm of the amounts used of target nucleic acid and determining a continuously differentiable function of the determined C_p values as a function of the logarithm of the amounts used of reference nucleic acid;
 - f) determination of the C_p values of the target nucleic acid and reference nucleic acid in the sample to be analysed as well as in a calibrator sample;
 - g) assignment of the C_p values measured in step f) to a particular function values of the functions determined in step e);
 - h) calculating the quotients of the function values from g) of the target nucleic acid and reference nucleic acid for the sample to analysed as well as for the calibrator sample; and
 - i) determination of the ratio of the two quotients from h) as a measure for the original amount of target nucleic acid contained in the sample.

10. (Amended) A method for relative quantification of a target nucleic acid relative to a reference nucleic acid and standardized with a calibrator sample comprising the steps of:
- a) preparation of a common or two separate dilution series of target nucleic acid and reference nucleic acid;
 - b) amplification of the various dilutions of target nucleic acid and reference nucleic acid under defined reaction conditions, the amplification of the nucleic acid being measured in real-time;
 - c) setting defined signal threshold values for the target nucleic acid and reference nucleic acid;

- Sub B2
- d) determining the cycle number Cp at which the signal threshold values defined for the target nucleic acid and reference nucleic acid are exceeded in each dilution;
- e) determining a continuously differentiable function of the logarithm of the amounts used of target nucleic acid as a function of the Cp values determined in d) and determining a continuously differentiable function of the logarithm of the amounts used of reference nucleic acid as a function of the determined Cp values;
- f) determining the Cp values of the target nucleic acid and reference nucleic acid in the sample to be analysed as well as in a calibrator sample;
- g) assignment of the Cp values measured in step f) to particular function values of the functions determined in step e);
- h) calculating the quotients of the function values from g) of the target nucleic acid and reference nucleic acid for the sample to analysed as well as for the calibrator sample; and
- i) determination of the ratio of the two quotients from h) as a measure for the original amount of target nucleic acid contained in the sample.
- AH

- Sub B3
11. (Amended) The method of claim 10, wherein the continuously differentiable functions from step e) are determined with a polynomial fit.
12. (Amended) The method of claim 10, wherein the amplified nucleic acids are detected with at least one fluorescent-labelled hybridization probe.
13. (Amended) The method of claim 12, wherein the amplified nucleic acids are detected with of FRET hybridization probes, molecular beacons, or TaqMan® probes.
14. (Amended) The method of claim 10, wherein the amplified nucleic acids are detected with a DNA-binding dye.

REMARKS

Claims 1-14 were pending. Applicants have amended claims 1-3, 5, 6 and 8-14, without narrowing their scope, hereinabove. A marked-up version of the claims with changes